

Common Structures in 2,3 and 4D Simplicial Quantum Gravity *

H.S.Egawa ^{a, b}, N.Tsuda ^{b †} and T.Yukawa ^{c, b}

^aDepartment of Physics, Tokai University Hiratsuka, Kanagawa 259-12, Japan

^bTheory Division, Institute of Particle and Nuclear Studies, KEK, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305, Japan

^cCoordination Center for Research and Education, The Graduate University for Advanced Studies, Hayama-cho, Miuragun, Kanagawa 240-01, Japan

Two kinds of statistical properties of dynamical-triangulated manifolds (DT mfd) have been investigated. First, the surfaces appearing on the boundaries of 3D DT mfd were investigated. The string-susceptibility exponent of the boundary surfaces ($\tilde{\gamma}_{st}$) of 3D DT mfd with S^3 topology near to the critical point was obtained by means of a MINBU (minimum neck baby universes) analysis; actually, we obtained $\tilde{\gamma}_{st} \approx -0.5$. Second, 3 and 4D DT mfd were also investigated by determining the string-susceptibility exponent near to the critical point from measuring the MINBU distributions. As a result, we found a similar behavior of the MINBU distributions in 3 and 4D DT mfd, and obtained $\gamma_{st}^{(3)} \approx \gamma_{st}^{(4)} \approx 0$. The existence of common structures in simplicial quantum gravity is also discussed.

1. Introduction

The development of simplicial gravity started with the 2D case, and has now reached the point of simulating the 4D case. The scaling structures of boundaries in 2,3 and 4D Euclidian space-time have been investigated [1–3]. Concerning the analogy of the loop-length distribution in 2D [1], the scaling relations of the boundary surface-area distribution in 3D and the boundary-volume distribution in 4D have been shown near to the critical point [2,3].

First, we are concerned with the surfaces appearing on boundaries in 3D DT mfd with S^3 topology. We actually investigated the relation between the boundary surfaces of 3D simplicial quantum gravity (SQG) and the random surfaces of 2D SQG. In order to investigate the statistical quantities of these boundary surfaces, we measure the string-susceptibility exponent ($\tilde{\gamma}_{st}^{(2)}$) of the boundary surface of 3D DT mfd by means of a MINBU[4] analysis. We denote $\tilde{\gamma}_{st}^{(d)}$ as the string-




Figure 1. Schematic picture of an analysis of the boundaries in 3 and 4D DT mfd with γ_{st} and $\tilde{\gamma}_{st}$.

susceptibility exponent of the boundary surfaces in $(d + 1)$ D DT mfd. Fig.1 shows a schematic picture of the boundary surfaces at a distance d in 3 and 4D DT mfd. We naively expected equivalence between the boundary surfaces of 3D DT mfd and the random surfaces on 2D DT mfd.

Second, we investigated the structures of 3 and 4D DT mfd [5] by measuring the stan-

*Presented by H.S. Egawa

[†]supported by Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists.

standard MINBU distributions, and obtain the string-susceptibility exponents ($\gamma_{st}^{(3)}$ and $\gamma_{st}^{(4)}$). We found the fact that $\gamma_{st}^{(3)} \approx \gamma_{st}^{(4)} \approx 0$. The aim of this short paper is to discuss the existence of common structures in simplicial quantum gravity.

2. Models

We start with the Euclidean Einstein-Hilbert action in d -dimensional SQG,

$$S_{EH}[\Lambda, G] = \int d^d \xi \sqrt{g} (\Lambda - \frac{1}{G} R), \quad (1)$$

where Λ is the cosmological constant and G is Newton's constant. We use the lattice action,

$$S_{Lattice}^{(d)}[\kappa_{d-2}, \kappa_d] = \kappa_d N_d - \kappa_{d-2} N_{d-2}, \quad (2)$$

where κ_{d-2} and κ_d are coupling constants. N_i denotes the number of i -simplexes. The lattice actions in 2,3 and 4D SQG are as follows:

$$S^{(2)}[\kappa_2] = \kappa_2 N_2 \quad (2D), \quad (3)$$

$$S^{(3)}[\kappa_3, \kappa_1] = \kappa_3 N_3 - \kappa_1 N_1 \quad (3D), \quad (4)$$

$$S^{(4)}[\kappa_4, \kappa_2] = \kappa_4 N_4 - \kappa_2 N_2 \quad (4D). \quad (5)$$

We consider a partition function,

$$Z(\kappa_{d-2}, \kappa_d) = \sum_{T(d)} e^{-S(\kappa_{d-2}, \kappa_d)}. \quad (6)$$

We sum over all simplicial triangulations ($T(d)$) in order to carry out a path integral about the metric.

3. Analysis of Boundaries

We now consider the boundary surfaces of 3D DT mfd with S^3 topology. The boundary surfaces are located at a distance d from the origin. The mother boundary surface is defined as the surface with the largest tip volume[6]. It is reported that the non-trivial scaling behavior of the surface area distribution of the mother boundary surface in 3D SQG has been shown to exist near to the critical point[2]. Therefore, we concentrate on the mother boundary surface for the measurements.

3.1. String-Susceptibility Exponent of the Mother Boundary in 3D SQG

We now show how to measure $\tilde{\gamma}_{st}$ of boundary surfaces having different sizes. Suppose that the spherical boundary surfaces are obtained; we can then use the MINBU algorithm to measure $\tilde{\gamma}_{st}$. In the case that the topology of the boundary surface is not spherical, but is a handle-body, such as a torus, the standard MINBU algorithm does not work well. Therefore, we have concluded that we should concentrate on spherical boundary surfaces. Fig.2 shows a measurement of the mother spherical boundary MINBU distributions of 3D DT mfd with $N_3 = 16K$. The lower limit of the volume of the boundary surfaces is 400. Our numerical result shows $\tilde{\gamma}_{st}^{(2)} \approx -0.5$. In 2D SQG with S^2 topology, the exact calculation of the string-susceptibility exponent ($\gamma_{st}^{(2)}$) is well-known to be $\gamma_{st}^{(2)} = -0.5$. We find a correspondence between the string susceptibility of the mother spherical boundary surfaces in 3D DT mfd and that of 2D DT mfd. Therefore, we have obtained evidence of an equivalence between the boundary surfaces of 3D SQG and the random surfaces of 2D SQG. When these boundary surfaces can be recognized as the surfaces of the matrix model, 3D DT mfd can be reconstructed by direct products of random surfaces of 2D QG and the geodesic distance.

4. String-Susceptibility Exponents in 3 and 4D SQG

The standard MINBU algorithms are not available for the case of the handle-body in an arbitrary dimension. It is not easy to choose 3D spherical boundary mfd in 4D DT mfd, because we cannot use the Euler's characteristic in 3D in order to distinguish the topologies of 3D DT mfd. It is therefore difficult to determine $\tilde{\gamma}_{st}^{(3)}$ of the boundary mfd in 4D DT mfd by means of the MINBU technique due to the reason mentioned above. However, we can compare it with the string-susceptibility exponents in 3 and 4D DT mfd. Fig.3 shows the MINBU distributions of 3D DT mfd with $N_3 = 16K$ near to the critical point ($\kappa_0 = 4.09$ and $\kappa_3 = 2.20$). We then




Figure 2. Boundary MINBU distributions in terms of the spherical boundary surfaces in 3D DT mfd at a distance of 10 with $N_3 = 16K$ using $\log - \log$ scales. The dots represent the raw-data points.

obtained $\gamma_{st}^{(3)} \approx 0$ of 3D DT mfd by a MINBU analysis. Moreover, the standard Minbu distributions of 4D DT mfd are measured near to the critical point; we actually obtained the string-susceptibility exponent as $\gamma_{st}^{(4)} \approx 0$.

5. Summary and Discussions

The space-time structures in 2 and 3D SQG were investigated using a boundary analysis by means of a MINBU analysis. We have obtained evidence of an equivalence between the boundary surfaces of 3D SQG and the random surfaces of 2D SQG with numerical results, $\tilde{\gamma}_{st}^{(2)} \simeq \gamma_{st}^{(2)}$. Furthermore, we obtained the string-susceptibility exponents: both $\gamma_{st}^{(3)} \approx 0$ in 3D SQG near to the critical point (belonging to the strong-coupling phase) and $\gamma_{st}^{(4)} \approx 0$ in 4D SQG near to the critical point (belonging to the strong-coupling phase). In the 3D case, we conjecture the relation $\tilde{\gamma}_{st}^{(3)} \simeq \gamma_{st}^{(3)}$, where $\tilde{\gamma}_{st}^{(3)}$ is obtained from the boundary mfd in 4D DT mfd. If the above conjecture is correct, the higher dimensional SQG (i.e., 3 and 4D SQG) can be reduced to the lower dimensional SQG (maybe 2D SQG). Some similar discussions are found in ref.7.




Figure 3. Standard MINBU distributions in 3D DT mfd with $N_3 = 16K$ using $\log - \log$ scales.

REFERENCES

1. H.Kawai, N.Kawamoto, T.Mogami, and Y.Watabiki, Phys.Lett. B306 (1993) 19; J.Ambjørn, J.Jurkiewicz and Y.Watabiki, Nucl.Phys. B454 (1995) 313; N.Tsuda and T.Yukawa, Phys.Lett. B305 (1993) 223.
2. H.Hagura, N.Tsuda and T.Yukawa, *Phases and fractal structures of three-dimensional simplicial gravity*, hep-lat/9512016.
3. H.S.Egawa, T.Hotta, T.Izubuchi, N.Tsuda and T.Yukawa, Prog.Theor.Phys. 97 (1997) 539; Nucl.Phys.B (Proc.Suppl.) 53 (1997) 760.
4. S.Jain and S.D.Mathur, Phys.Lett.B286 (1992) 236.
5. J.Ambjørn, S.Jain, J.Jurkiewicz and C.F.Kristjansen, Phys.Lett. B305 (1993) 208; S.Catterall, G.Thorleifsson, J.Kogut and R.Renken Nucl.Phys. B468 (1996) 263.
6. H.S.Egawa and N.Tsuda, these proceedings.
7. I.Antoniadis, P.O.Mazur and E.Mottola, Phys.Lett. B394 (1997) 49.